

Title

Can the collective cognitions of individual professionals within healthcare teams predict the team's performance: developing methods and theory.

Authors

Martin P Eccles [1] martin.eccles@ncl.ac.uk

Susan Hrisos [1] Susan.hrisos@ncl.ac.uk

Jillian J Francis [2] j.francis@abdn.ac.uk

Nick Steen [1] nick.steen@ncl.ac.uk

Marie Johnston [3] m.johnston@abdn.ac.uk

Institutions

1. Institute of Health and Society, Newcastle University
2. Health Services Research Unit, University of Aberdeen
3. Department of Psychology, University of Aberdeen

Corresponding Author:

Professor Martin Eccles, Institute of Health and Society, University of Newcastle upon Tyne,

21 Claremont Place, Newcastle upon Tyne NE2 4AA

Phone 0191 2228674, Fax 0191 2226043, email martin.eccles@ncl.ac.uk

Abstract

Background

Within Implementation Research, it can be useful to use theory based approaches to understanding the behaviours of healthcare professionals, the quality of care that they reflect and to design interventions to change them. However, such approaches lead to a new range of methodological and theoretical challenges including how to measure and appropriately relate the predictors of behaviour and the behaviours of healthcare professionals working in teams. The aim of this study was to explore the relationship between the Theory of Planned Behaviour proximal predictors of behaviour (intention and perceived behavioural control) and practice level behaviour. This was done in the context of two clinical behaviours (statin prescription and foot examination) in the management of patients with diabetes mellitus in primary care. Scores for the predictor variables were aggregated over healthcare professionals using four methods: (1) simple mean of all primary care team members' intention scores; (2) highest intention score combined with perceived behavioural control (PBC) of the highest intender in the team; (3) highest intention score combined with the highest PBC score in the team; (4) the scores (on both constructs) of the team member identified as having primary responsibility for the clinical behaviour.

Methods

Scores on theory based cognitive variables were collected by postal questionnaire survey from a sample of primary care doctors and nurses from north east England and the Netherlands. Data on two clinical behaviours were patient reported and collected by postal questionnaire survey. Planned analyses explored the predictive value of various aggregations of intention and perceived behavioural control in explaining variance in the behavioural data.

Results

Across the two countries and two behaviours, responses were received from 37-78% of healthcare professionals in 57-93% practices; 51% (UK) and 69% (Netherlands) of patients surveyed responded. None of the aggregations of cognitions predicted statin prescription. The highest intention in the team (irrespective of PBC) was a significant predictor of foot examination.

Conclusions

These approaches to aggregating individually-administered measures may be a methodological advance of theoretical importance. Using simple means of individual-level measures to explain team-level behaviours is neither theoretically plausible nor empirically supported; the highest intention was both predictive and plausible. In studies aiming to understand the behaviours of teams of healthcare professionals in managing chronic diseases some sort of aggregation of measures from individuals is necessary. This is not simply a methodological point but a necessary step in advancing the theoretical and practical understanding of the processes that lead to implementation of clinical behaviours within healthcare teams.

Background

Within Implementation Research - the scientific study of methods to promote the uptake of research findings, and hence to reduce inappropriate care – using theory based approaches to understanding the behaviours of healthcare professionals and the quality of care that they reflect and designing interventions to change them is being promoted [1, 2]. However, such approaches lead to a new range of methodological and theoretical challenges pre-eminent amongst which are how to measure and appropriately relate predictors of behaviour and the behaviour of healthcare professionals [3].

It is possible, and in certain circumstances appropriate and feasible, to directly observe the behaviour(s) of healthcare professionals (HCPs). However, this is likely to be expensive, time consuming and ethically problematic. For studies that involve large numbers of HCPs other methods of measuring HCP behaviour using various forms of routinely available data can be more economical. Such data that reflect a proxy, or indirect, measure of HCP behaviour usually fall into two categories; recorded measures of HCP behaviour and measures (clinical, physiological or biochemical) of the patient's condition. An example of the former would be the prescription of a statin reflecting behaviour in relation to the management of hypercholesterolaemia. An example of the latter would be a patient's serum cholesterol level. Each of these has its own threats to its validity as a measure of an individual HCPs behaviour.

Commonly (at least within the UK and the Netherlands), data collected will indicate that the clinical behaviour has been done, but then it may not be possible to know who within the clinical team did it. Thus the threat to the use of recorded data is that the measure may not relate uniquely to a single HCP, but to the collective actions of the team. Prescriptions attributed to a doctor may have been issued by trainee doctors or locums plus the prescribed treatment of an individual patient may be changed by different doctors within a practice over time. Similarly, for clinical measures of the patient's condition (setting aside considerations of patient adherence and physiology), a threat to validity would be that a measure of a patient's serum cholesterol may also reflect the behaviours of more than one HCP. A nurse may advise a patient about their diet and a doctor may prescribe a statin. Although illustrated with a simple example these issues apply to any chronic condition managed by a team of

healthcare professionals in primary care e.g. diabetes, heart disease, asthma or chronic obstructive airways disease. Such threats to validity can obscure attempts to explain the behaviour of HCPs. It is therefore important to develop methods of predicting clinical behaviours that can take account of the collective performance of individuals working in teams.

Theoretical context

Explanations for clinical behaviour can be investigated using psychological theories which have been successful in predicting behaviour and behaviour change in other settings. Using such a theory based approach offers the potential of a generalisable framework within which to consider factors influencing behaviour and the development of interventions to modify them. A study by Eccles et al [3] used six theories to investigate factors associated with prescribing antibiotics for patients with a sore throat amongst primary care doctors. This showed that the impact of individual beliefs and perceptions on intention to prescribe was high, including both evidence-based and non-evidence based factors, whilst the impact on behaviour was considerably smaller. Two systematic reviews of the relationship between intention and behaviour in individual HCPs [4, 5] found only 16 eligible studies but suggested that the nature of the relationship was similar to that shown by reviews of much larger numbers of studies in non-healthcare professionals [6]. Data such as these allow clear predictions to be made about the factors likely to change psychological constructs and to change behaviour.

One of the more widely used theories is the Theory of Planned Behaviour (TPB) [7]. The TPB proposes a model about how human action is guided. It predicts the occurrence of a specific behaviour provided that the behaviour is intentional (i.e. the model does not claim to predict behaviours that are habitual or automatic). The TPB model is shown in Figure 1 and depicts the three cognitive variables that the theory suggests will predict the intention to perform a behaviour. Whilst intention is the main precursor of behaviour, perceived behavioural control (PBC) also directly predicts behaviour. For example, a positive intention may be prevented from being translated into action because of an internal or external barrier that the individual perceives as insurmountable.

Because data may reflect the behaviour of more than one HCP, it is thus appropriate to analyse these proxy behaviour data at the aggregated level of a primary care practice. Thus, recorded data can indicate reliably that a patient has been prescribed a statin by one of the HCPs in the practice. However, in order to use a theory based approach it is then necessary to also consider aggregating individuals' measurement of cognitions (about the prescribing of statins). It would be possible to aggregate measures of individuals' cognitions about clinical behaviours and conditions as a simple mean (as is the practice in the literature on measurement of team level variables such as team climate[8]). However, the mean may not reflect the organisational, professional and social processes involved in the team. It may be possible to improve the predictive performance of measures that represent team cognitions by taking account of factors such as individuals' roles, responsibilities or positions. For example, when identified individuals predominantly perform, or have responsibility for, a behaviour (foot examination of patients with diabetes by a practice nurse) then that individual's intention score could be used as a sole representative measure or used to weight a mean value.

Clinical context

Type 2 diabetes mellitus (DM) is an increasingly prevalent chronic illness and is an important cause of avoidable mortality. Studies of the quality of care for patients with diabetes suggest less than optimum care in a number of areas [9]. In primary care the management of DM includes glycaemic control; blood pressure control; foot examination for peripheral pulses and neuropathy; lipid control; and weight reduction (retinopathy screening is often organised separately from the practice). Patients are managed by the integrated activities of medical and non-medical members of the primary care team.

Aim

The aim of this study was to explore the relationship between the Theory of Planned Behaviour's direct predictors of behaviour (individuals' intention and perceived behavioural control) aggregated over HCPs in a number of ways, and practice level behaviour in the context of care for patients with diabetes mellitus in primary care.

The method of aggregation is not simply a statistical device but may reflect different team processes and different theoretical approaches to team-functioning. For example,

aggregating intentions by averaging suggests equal weighting of members' views and would suggest team decision-making based on equal and shared communications. Whereas, choosing the highest intention score in the team to represent the relevant 'team cognition score', suggests that the team has allocated roles, with one member specialising in, or having responsibility for, the targeted clinical behaviour; here the underlying model suggests a more complex team structure with more stream-lined decision-making. Other methods of aggregating would also test specific role structures e.g. the team process may be best assessed by selecting the highest intention indicating responsibility for decision-making along with highest PBC indicating responsibility and capability for the actual behaviour. Therefore we investigated the following methods of aggregating respondents' scores for each primary care team: (1) simple mean of all PC team members' intention scores; (2) highest intention score from responding HCPs combined with PBC for either that individual, or (3) for the highest PBC (4) the scores of the HCP identified as having primary responsibility for the clinical behaviour, ignoring the scores of other team members.

Methods

Design and participants

This was a predictive study of the theory based cognitions and clinical behaviours concerning the management of patients with diabetes of a sample of primary care doctors and nurses from north east England and the Netherlands. Data on cognitions was collected by postal questionnaire survey; behavioural data was patient reported and collected by postal questionnaire survey. Planned analyses explored the predictive value of various aggregations of intention and perceived behavioural control in explaining variance in the behavioural data.

Study setting

The study was based within two randomised controlled trials of interventions to improve the management of patients with diabetes cared for in primary care.

Study practices

In the UK the study practices were those in three Primary Care Trusts (PCTs) served by two district hospital based diabetes registers both using the same register software. In the Netherlands the practices were those in three regions of the middle and south of the Netherlands [10].

Study patients

In the UK the study patients were those people with type 2 diabetes appearing on the area wide diabetes registers, aged over 35 and receiving diabetes care exclusively from the DREAM trial (The **D**ialysis **R**Ecall **A**nd **M**anagement system trial) [11] practices or shared between study practices and hospital. In the Netherlands patient reported outcomes were gathered from patients with type 2 diabetes, who were younger than 80 years and registered with practices participating in the PAS trial (The diabetes **P**assport as an **A**id to **S**tructure diabetes management in primary care trial) [10]. Patients managed in secondary care were excluded from the PAS trial.

Predictive measures

Theoretically derived measures were developed following the operationalisation protocols of Ajzen [7, 12]. Twelve UK primary care doctors and practice nurses were interviewed about three behaviours (measuring blood pressure, foot examination, prescribing statins). The

schedule for these semi-structured interviews was designed to elicit responders' beliefs relating to the constructs of the TPB. Primary care doctors and practice nurses were encouraged to talk freely about these beliefs and any ambiguities were clarified using appropriate prompts. Interviews were tape recorded, transcribed and content analysed. Beliefs frequently mentioned in the interviews were used to design items in a questionnaire that was developed for each of the three behaviours. The response format for all items was a seven point Likert-type scale, from 1 (strongly agree) to 7 (strongly disagree). This initial draft of the questionnaire was pre-tested with a further six UK primary care doctors for style and clarity of content and to determine completion time. Minor revisions of wording were made to the questionnaire in the light of their comments. The final questionnaire used in the UK covered three behaviours, both 'indirect' and 'direct' measures of the theoretical constructs [7, 12] and consisted of 154 items, including questions about the size of practices and demographic details. For the Netherlands survey because of concerns about respondent burden, a shortened set of the questions from the UK questionnaire was used covering only two of the three behaviours and using only direct measures. The relevant questions from the UK set were translated into Dutch and then back translated into English (and adjusted where necessary) to ensure that the meaning was the same for the UK and Dutch studies. The questions measuring intention and perceived behavioural control for the two behaviours of prescribing statins and examining patients' feet are shown in Table 1. Scoring was adjusted so that a high score indicates a strong intention and a high degree of perceived control.

Outcome measures

In the UK, as part of a larger patient reported outcomes survey [11], patients with type 2 diabetes were asked the following two questions. Firstly, "Please provide as much information as you can in the box below about **ALL** the medication you have taken **over the last four weeks**"; any report of a statin was identified. Secondly, they were asked, "Over the last 12 months did you have any of the tests or investigations listed"; the list included: "test of feeling on your feet"; a positive response was taken as an indication of having a foot examination. In the Netherlands patients were asked to report on the medication they were currently taking and whether or not they had had their feet examined in the past 15 months.

For both countries, responses were used to calculate the percentage of patients per practice who reported taking a statin and the percentage of patients per practice who reported having their feet examined.

Procedure

In both the UK and the Netherlands the questionnaire was mailed to all primary care doctors and nurses at participating trial practices at the end of the intervention period. In the UK two reminder letters were sent to non responders at fortnightly intervals. Dutch non-responders received one reminder letter after three weeks. Patient reported outcomes were also collected by postal questionnaire at the end of the intervention period of both trials.

Analytical approach

Internal consistency of multi-item measures [of intention and PBC] was assessed using Cronbach's alpha (for measures with >2 items) using an acceptability criterion of $\alpha > 0.6$, and Pearson's correlation coefficient (for 2 item measures) using an acceptability criterion of $r > 0.25$.

We were interested in the relationship between practice level behaviour and aggregations of individuals' cognitions (intentions and perceived behavioural control) and investigated this using multiple regression analysis. We conducted analyses to reflect four possible team patterns. First, we argued that the behaviour was likely to be driven equally by the individual intentions of all the practice members; we therefore calculated a mean value for each practice. It was likely that we would both get responses from single doctor practices and get single responses (from either a nurse or a doctor) from multi-doctor practices. Under these circumstances the concept of a mean value was less meaningful and therefore, we repeated the analyses including only those practices from which we received more than one response. Second, we considered that behaviour could be most driven by the individual with the highest intention (and their PBC) within the practice and so used these measures as predictor variables. Third, we considered that the behaviour could be the product of one team member having a strong intention and another team member having a high level of PBC. An example of this would be the situation where a nurse had a high intention to perform the behaviour and a doctor had a high PBC score as a consequence of knowing that the nurse intended to perform the behaviour. Fourth, we considered that behaviour was most likely to be driven by

the individual whose role it was to perform the behaviour. Therefore, for foot examination, we considered that this could be the role of a nurse. The stratified analysis was restricted to doctors.

As the Theory of Planned Behaviour predicts a direct effect of both intention and perceived behavioural control (PBC) on behaviour, both were included in the regression analyses. We also explored a country effect (to allow for both 'real' and methodological differences between them) and the number of responses per practice. Although both host studies were randomised controlled trials we analysed them as two cross sectional studies on the basis that any effect of the interventions on behaviour would be mirrored by a change in cognitions and that the relationship between cognitions and behaviour should therefore persist, whether or not the trial changed the levels observed in the intervention group.

Ethical approval

The UK study was approved by the South Tyneside, Southwest Durham, Hartlepool and North Tees Local Research Ethics Committees (LRECs). The Dutch study was approved by the ethics committee of Radboud University Medical centre, Nijmegen, The Netherlands.

Results

The details of the number of healthcare professionals surveyed and the characteristics of their practices and the survey response rates are shown in Table 2. Practices were dichotomised into single or multi-practitioner practices. The 69 practices contributing at least one responder to the analysis were not significantly different in terms of size to non-responder practices (Pearson $\chi^2 = 2.248$, d.f. = 1, $p = 0.13$). For the analysis of foot examination, the number of nurses per practice was also available. Practices were again dichotomised and the 83 practices contributing at least one responder to the analysis were not significantly different in terms of the number of primary care doctors in the practice (Pearson $\chi^2 = 2.149$, d.f. = 1, $p = 0.14$); but were significantly more likely to have two or more nurses (80% v 47%, Pearson $\chi^2 = 7.215$, d.f. = 1, $p = 0.007$).

In the UK study a random sample of 2815 patients were surveyed and usable responses were received from 1433 (51%). In the Dutch study 1432 patients were surveyed with 993 (69%) usable responses received. Overall, 736/2426 (30%) patients reported taking statins (362/1433 (25%) UK patients and 374/993 (38%) Dutch patients). Overall, 1234/2426 (51%) patients reported having their feet examined in the past 12 (UK) or 15 (Dutch) months (806/1395 (58%) UK patients and 428/993 (43%) Dutch patients).

Prescribing statins

The three item measure of intention had a Cronbach's alpha of 0.95. The two item measure of PBC had a Pearson's Correlation Coefficient of 0.37 ($p < 0.001$). In UK practices the overall mean (sd) of the practice mean intention score was 4.8 (1.5) and in Dutch practices this was 5.6 (1.3) (mean difference (95% CI) -0.7300 (-1.4 to -0.04) $p = 0.038$). Similar values for the strongest intention were, for the UK practices, 5.2 (1.5) and for the Dutch practices 5.7 (1.3); these were not significantly different. The mean intention score (from participating HCPs) within each practice was significantly correlated with the highest intention score within that practice (Pearson Correlation Coefficient 0.93, $p < 0.001$) but neither was significantly correlated with the practice mean percentage of patients taking a statin.

In a regression model including both mean intention and mean PBC (Table 3), neither significantly predicted behaviour but there was a significant "country effect" with Dutch primary care doctors being 11% more likely to prescribe statins. When PBC was removed

from the model intention still did not predict behaviour and there was no additional effect of an interaction term between intention and country (i.e. intention was not a significantly greater predictor in one country than the other). A similar analysis restricted to the smaller number of practices where there was more than one respondent produced a similar pattern of results though the country effect was not significant.

When using the highest intention score for each practice, none of highest intention, PBC of the highest intender or highest PBC in the practice predicted the prescription of statins (Table 3). Again the country effect is apparent and of the same order of magnitude and significance. When PBC was removed from the model intention still did not predict behaviour and there was no additional effect of an interaction term between intention and country.

Foot examination

The three item measure of intention had a Cronbach's alpha of 0.96. The two item measure of PBC had a Pearson's Correlation Coefficient of 0.44 ($p < 0.001$). In UK practices the overall mean (sd) of the practice mean intention score was 4.9 (1.3) and in Dutch practices this was 4.4 (1.4); these were not significantly different. Similar values for the strongest practice intention were, for the UK practices, 5.9 (1.3) and for the Dutch practices 5.1 (1.6) (Mean difference (95%CI) 0.78 (0.14 to 1.43), $p = 0.018$). The mean intention score for a practice was significantly correlated with the highest intention score within that practice (Pearson Correlation Coefficient 0.78, $p < 0.01$) and the highest intention score was also significantly correlated with the practice mean percentage of patients reporting a foot examination (Pearson Correlation Coefficient 0.29, $p < 0.01$).

In a regression model (Table 3) including both mean intention and mean PBC, neither significantly predicted behaviour but there was a significant "country effect" with UK practices being 14% more likely to inspect feet. When PBC was removed from the model intention still did not predict behaviour and there was no additional effect of an interaction term between intention and country. A similar analysis restricted to the smaller number of practices where there was more than one respondent produced a similar pattern of results though the country effect was not significant.

The highest intention score in a practice belonged to 38 nurses (24 of whom were from practices where intention scores were available for both primary care doctor and nurse

respondents) and 39 primary care doctors (eight of whom were from practices where intention scores were available for both primary care doctor and nurse respondents). In the remaining six practices this score was the same for both nurse and primary care doctor and the regression used the scores for individuals who have both the highest intention and the highest PBC. The highest practice intention was a significant predictor of foot examination. Again, there was a significant country effect with reported foot inspections being 11% fewer in ND practices than UK practices ($p=0.011$). Removing PBC, including an interaction term for intention/country and including type of healthcare professional (thus exploring professional role) did not significantly change the model. Finally the analysis was repeated using the highest intention score for the practice and the strongest PBC score for the practice. In this model, the PBC score is predominantly that of the primary care doctor respondents. This analysis produced results similar to the previous one.

Discussion

This paper reports an analysis of four different ways of dealing with the problem of relating the cognitions of individual members of a team of healthcare professionals to a shared outcome of their collective behaviours. For the behaviour of foot examination, how the individual cognitions were analysed made a difference with strongest intention, not mean intention, being significantly associated with practice level behaviour. However, this has to be regarded as exploratory and preliminary in a number of ways.

The theories we were using were not necessarily intended to be used as we have used them and we are proposing an extension of the use of the TPB to the collective behaviour of a team. Pragmatically there does not seem to be any reason why measures cannot be used in this way. Indeed, other measures of team performance, such as the Team Climate Inventory, use a simple mean as their summary statistic [8]. In a theoretical context, it is unclear what a team's mean intention score represents. However, as suggested earlier, if mean intention is predictive it suggests some kinds of collective processes, especially with regard to decision making and communication. Our finding that mean intention was not predictive (whilst acknowledging our limited numbers and response rates), suggests that for the management of these two clinical behaviours by primary care teams, decision-making and responsibility may not be distributed equally across the team.

We were using a cognitive model for what seem to be intentional behaviours. However, these are relatively routine behaviours and they may well, over time, become routinely maintained and therefore no longer need thinking through each time they are performed. Therefore other measures, either instead of or alongside cognition models, may have additional predictive power for teams. Indeed, in a study of primary care doctor's practitioners' antibiotic prescribing behaviour that compared the predictive power of theories, a measure of habit was the best predictor of behaviour [3].

Whilst mean levels of intention to perform both behaviours were positive being between 4.4 and 5.6 for both behaviours in both countries, levels of performance for what should be almost universal behaviours were low; for only foot examination in the UK was the reported rate of performance about 50%. This could be due to: low reporting rates by patients (our source of this data); the potential mismatch for prescribing statins engendered by patients

reporting what they were taking and doctors reporting their intention to prescribe; or over estimation of intention by healthcare professionals. This raises the possibility of there being post-intentional factors that are influencing behaviour. Such factors include situation-behaviour associations that may become salient as a result of planning interventions. There is evidence that other relevant modifiers of the intention-behaviour relationship are intention certainty, past behaviour, self-schema and anticipated regret. In addition, whether the behaviour is more likely to be influenced by internal factors (e.g., attitudes) or by external factors (e.g., perceived normative constraints) and temporal stability are potential influences [13]. The fact that we had no explicit measurement of such factors may have limited our ability to explain variation in behaviour. However, there is a limit to how many theoretical constructs can be included and operationalised within a single instrument and we already had problems with response rates that could have been a function of questionnaire length.

The finding that the strongest intention score within each team, for inspecting feet, significantly predicted patients' reports of foot inspection, is consistent with the possibility that healthcare professionals may have had stronger intentions if they had been assigned responsibility for foot inspection within the practice (though our attempt to allocate roles in our analyses did not confirm this). The idea that assigned roles and responsibilities influence cognitions and behaviour has received substantial support in the behavioural literature [14, 15]. An alternative possibility is that teams allocate responsibility for a task to those with the strongest intentions to perform it, i.e. that roles evolve and may be chosen rather than being allocated. These possibilities warrant further investigation.

Whilst we explored different ways of relating behaviour and its theorised predictors our data from healthcare professionals had limitations. We know that across individual practices we usually had only a minority of team members responding so that our team mean values did not include values from those disinclined to complete questionnaires. The implication of this is that we may have lacked the power to detect difference across our different analyses. Also, if individual healthcare professionals do have a specified role within a practice (e.g. to inspect patients' feet), we do not know whether that individual responded to the questionnaire. If individuals with the highest intention within the team, or with the assigned responsibility, did not respond, then we may have underestimated these effects. Whilst non-response is an

enduring issue for health services research in general an ideal study of this type would include responses from all members of the participating teams.

Conclusions

However exploratory this work, the issues raised are of enduring importance, both methodologically and theoretically. In studies wishing to understand the behaviours of healthcare professionals in relation to the management of many chronic diseases then some sort of aggregation of measures from individuals is inevitably going to be necessary. Given that so much of healthcare involves teams of healthcare professionals the issues addressed in this study, however imperfectly, need to be addressed. This is not simply a methodological point but a necessary step in advancing the theoretical and practical understanding of the processes that lead to implementation of clinical behaviours within healthcare teams.

Competing interests

The authors declare that they have no competing interests.

Authors contributions

MPE, MJ and JF conceived the study. MPE, JF and SH were responsible for data collection. MJ and NS supervised the analyses. MPE led the writing and all authors commented on sequential drafts and approved the final version of the manuscript.


Acknowledgements

We are grateful to the participants in the two studies that provided the data for the analyses reported in this paper. We are grateful to: Dr R Dijkstra, Dr J Braspenning and Prof R Grol for access to data from the PAS Trial; and to Marije Bosch for work on the Dutch postal survey.

Figure Legends

Figure 1. The Theory of Planned Behaviour [7].

Figure 1.



Note. The three proximal variables also influence one another. Although this figure is presented in a simplified form, a more detailed diagram would include double-ended arrows joining these three variables.

Table 1. Questions measuring intention and perceived behavioural control for the two clinical behaviours.

| |
|---|
| Each question in the following section refers to the PRESCRIBING OF STATINS to your patients with Type 2 diabetes |
| Intention questions |
| I intend to prescribe a statin to most of the patients I see in the next month |
| I expect to prescribe a statin to most of the patients I see in the next month |
| I want to prescribe a statin to most of the patients I see in the next month |
| Perceived behavioural control questions |
| To prescribe a statin is easy |
| Overall, I feel that I can prescribe statins if I want to |
| Each of the questions in the following section refers to FOOT EXAMINATIONS on your patients with Type 2 diabetes |
| Intention questions |
| I intend to examine the feet of all my patients I see in the next month who have not been examined by the chiropodist or the podiatrist |
| I expect to examine the feet of all my patients I see in the next month who have not been examined by the chiropodist or the podiatrist |
| I want to examine the feet of all my patients I see in the next month who have not been examined by the chiropodist or the podiatrist |
| Perceived behavioural control questions |
| Examining patients' feet is easy |
| Overall I feel that I can examine these patients' feet if I want to |

Table 2. Characteristics of sample and questionnaire response rates from healthcare professionals for the two behaviours.

| | Overall | | | Response rates (n(%)) | | | | | |
|---|---------|---------|---------|-----------------------|---------|----------|------------------|---------|----------|
| | Numbers | | | Statin prescription | | | Foot examination | | |
| | UK | Dutch | Total | UK | Dutch | Total | UK | Dutch | Total |
| Number of HCPs | | | | | | | | | |
| primary care doctors | 161 | 59 | 220 | 59 (37) | 46 (78) | 105 (48) | 59 (37) | 46 (78) | 105 (48) |
| Nurses | 119 | 22* | 141 | | | | 53 (45) | 19 (86) | 72 (51) |
| Practices | | | | | | | | | |
| Overall | 58 | 40 | 98 | 34 (57) | 35 (88) | 69 (70) | 46 (79) | 37 (93) | 83 (85) |
| Single primary care doctor | 15 | 15 | 30 | | | | | | |
| >1 primary care doctor | 43 | 25 | 68 | | | | | | |
| Number (Median (range))/practice | | | | | | | | | |
| primary care doctors | 2 (1-9) | 2 (1-4) | 2 (1-9) | | | | | | |
| Nurses | 2 (1-6) | 2 (1-5) | 2 (1-6) | | | | | | |

*Includes 8 nurses and 14 assistants who inspect feet; excludes 26 assistants who did not inspect feet.

Table 3. Regression models for mean and strongest intention for statin use and foot examination.

| Model | n | Predictive Variables | B | Beta | R ² (adj) | F | p value |
|---|----|---------------------------|--------|----------|----------------------|-------|---------|
| Prescribing statins | | | | | | | |
| Mean intention (all practices) | 69 | Mean intention | 0.005 | 0.05 | | | |
| | | Mean PBC | -0.006 | -0.034 | | | |
| | | Country | 0.11 | 0.389** | 0.127 | 4.312 | 0.008 |
| Mean intention (practices with >1 respondent) | 25 | Mean intention | -0.003 | -0.036 | | | |
| | | Mean PBC | 0.044 | 0.237 | | | |
| | | Country | 0.093 | 0.350 | 0.03 | 1.239 | 0.321 |
| Highest Intention (a) | 69 | Highest intention | 0.001 | 0.011 | | | |
| | | PBC of strongest intender | 0.016 | 0.108 | | | |
| | | Country | 0.115 | 0.406*** | 0.136 | 4.560 | 0.006 |
| Highest Intention (b) | 69 | Highest intention | 0.003 | 0.027 | | | |
| | | Highest PBC | 0.001 | 0.006 | | | |
| | | Country | 0.113 | 0.401*** | 0.125 | 4.244 | 0.008 |
| Foot examination | | | | | | | |
| Mean intention (all practices) | 83 | Mean intention | -0.003 | -0.017 | | | |
| | | Mean PBC | -0.016 | -0.084 | | | |
| | | Country | -0.125 | -0.322** | 0.097 | 3.922 | 0.012 |
| Mean intention (practices with >1 respondent) | 51 | Mean intention | -0.001 | -0.006 | | | |
| | | Mean PBC | 0.000 | 0.000 | | | |
| | | Country | -0.075 | -0.224 | -0.11 | 0.826 | 0.486 |
| Highest Intention (a) | 83 | Highest intention | 0.033 | 0.229* | | | |
| | | PBC of Highest intender | -0.008 | -0.054 | | | |
| | | Country | -0.113 | -0.289* | 0.138 | 5.390 | 0.002 |
| Highest Intention (b) | 83 | Highest intention | 0.034 | 0.239* | | | |
| | | Highest PBC | -0.008 | -0.048 | | | |
| | | Country | -0.116 | -0.297** | 0.138 | 5.363 | 0.002 |

References

1. Eccles M, Grimshaw J, Walker A, Johnston M, Pitts N: **Changing the behaviour of healthcare professionals: the use of theory in promoting the uptake of research findings.** *J Clin Epidemiol* 2005, **58**:107-112.
2. The Improved Clinical Effectiveness through Behavioural Research Group (ICEBeRG): **Designing theoretically-informed implementation interventions.** *Implementation Science* 2006, **1**:4.
3. Eccles MP, Grimshaw J, Johnston M, Steen IN, Pitts NB, Thomas R: **Applying psychological theories to evidence-based clinical practice: Identifying factors predictive of managing upper respiratory tract infections without antibiotics.** *Implementation Science* 2007, **2**:26.
4. Eccles MP, Hrisos S, Francis J, Kaner E, Dickinson HO, Beyer F, Johnston M: **Do self-reported intentions predict clinicians' behaviour: a systematic review.** *Implementation Science* 2006, **1**:28.
5. Godin G, Belanger-Gravel A, Eccles M, Grimshaw J: **Healthcare professionals' intentions and behaviours: A systematic review of studies based on social cognitive theories.** *Implementation Science* 2008, **3**(36).
6. Conner M, Armitage CJ: **Extending the theory of planned behavior: a review and avenues for further research.** *J Appl Psychol* 1998, **28**(15):1429-1464.
7. Ajzen I: **The theory of planned behaviour.** *Organizational Behaviour and Human Decision Processes* 1991, **50**:179-211.
8. West MA, Wallace M: **Innovation in health care teams.** *Eur J Soc Psychol* 1991, **21**(4):303-315.
9. Seddon ME, Marshall MN, Campbell SM, Roland MO: **Systematic review of studies of quality of clinical care in general practice in the UK, Australia and New Zealand.** *QHC* 2001, **10**(3):152-158.
10. Dijkstra R, Braspenning J, Grol R: **Implementing diabetes passports to focus practice reorganization on improving diabetes care.** *International Journal of Quality in Health Care* 2008, **20**:72-77.
11. Eccles MP, Whitty PM, Speed C, Steen IN, Vanoli A, Hawthorne GC, Grimshaw JM, Wood LJ, McDowell D: **A pragmatic cluster randomised controlled trial of a Diabetes REcall And Management system: the DREAM Trial.** *Implementation Science* 2007, **2**:6.
12. Francis J, Eccles MP, Johnston M, Walker AE, Grimshaw JM, Foy R, Kaner EFS, Smith L, Bonetti D: *Constructing questionnaires based on the theory of planned behaviour. A manual for health services researchers.* Newcastle upon Tyne: Centre of Health Services Research, University of Newcastle upon Tyne; 2004.
13. Sheeran P, Abraham C: **Mediator of moderators: temporal stability of intention and the intention-behavior relation.** *Society for Personality and Social Psychology* 2003, **29**(2):205-215.
14. Biddle BJ: *Role Theory--Expectations, Identities, and Behaviors.* New York: Academic Press; 1979.
15. Hardy ME, Conway ME: *Role theory: Perspectives for health professionals.* 2nd edition. Norwalk, CT: Appleton & Lange-Century-Crofts; 1988.